

# Evaluation of the periodontal status of a patient diagnosed with Sheehan syndrome: A case report

Seval Ceylan Şen<sup>1</sup>, Nuriye Işıl Saygun<sup>1</sup>, Arzu Or Koca<sup>2</sup>, Erkan Özcan<sup>1</sup>

<sup>1</sup> University of Health Sciences, Gülhane Faculty of Dental Medicine, Department of Periodontology, Ankara, Turkey

<sup>2</sup> University of Health Sciences, Ankara Keçiören Training and Research Hospital, Department of Endocrinology and Metabolic Diseases, Ankara, Turkey

## Abstract

**Aim:** Some immunological changes occurring in mothers during pregnancy increase their susceptibility to infections, including periodontal infections. Sheehan syndrome (SS) is a disease that occurs because of severe postpartum hemorrhage causing ischemic pituitary necrosis. The presence of multiple hormonal deficiencies leads to the impairment of bone microarchitecture, which can cause osteopenia and even osteoporosis. Osteoporosis and periodontitis are both chronic diseases characterized by bone loss. Moreover, recent studies have shown that there is a relationship between menopause, osteoporosis, alveolar bone resorption, and tooth loss. This case report aims to evaluate the oral and periodontal status of a patient with SS and to raise awareness about dental and periodontal problems that may occur in such patients.

**Methodology:** A 63-year-old female patient diagnosed with SS in the endocrinology clinic was referred to the periodontology clinic with complaints of multiple tooth loss, mobility in her teeth, and gingival bleeding. In the clinical and radiographic examination of the patient, it was determined that all teeth had horizontal bone resorption; moreover, there was mobility in the mandibular anterior teeth and hemorrhage in the gums. When the patient's laboratory results were examined, it was seen that the basal plasma levels of free triiodothyronine (T3) and thyroxine (T4), growth hormone (GH), prolactin (PRL), follicle stimulating hormone (FSH), luteinizing hormone (LH), progesterone, and total testosterone values were significantly lower, and the anterior pituitary elevation was significantly decreased in pituitary magnetic resonance (MR) imaging.

**Conclusion:** The periodontium is the target tissue of sex and other hormones. Hormonal changes may affect the inflammatory-immune tissue responses of periodontal tissues. Many studies have shown that thyroid and sex hormones affect oral and especially periodontal tissues. The degradation of hormonal hemostasis may induce a series of pathological events in the oral environment, resulting in inflammatory changes in gingival tissues, periodontal attachment losses, and destruction of the alveolar bone. Systemic diseases that affect hormonal conditions, such as SS, must be considered in the evaluation of oral health. Medical physicians should refer their patients to the dentist for consultation when evaluating patients' hormonal status and planning their treatment.

**Keywords:** Sheehan syndrome, hormones, periodontal status.

## Correspondence:

Dr. Seval CEYLAN ŞEN  
University of Health Sciences, Gülhane  
Faculty of Dental Medicine,  
Department of Periodontology, Etlik  
Keçiören, Ankara, Turkey  
E-mail: dt.sceylan@hotmail.com

Received: 6 March 2021

Accepted: 19 May 2021

## Access Online



DOI:

10.5577/intdentres.2021.vol11.suppl1.46

**How to cite this article:** Ceylan Şen S, Saygun NI, Or Koca A, Özcan E. Evaluation of the periodontal status of a patient diagnosed with Sheehan syndrome: A case report. Int Dent Res 2021;11(Suppl.1):308-11. <https://doi.org/10.5577/intdentres.2021.vol11.suppl1.46>

## Introduction

The periodontium is a tissue targeted by sex and other hormones. Indeed, hormonal changes may affect the inflammatory-immune tissue responses of periodontal tissues. Moreover, the degradation of hormonal hemostasis may induce a series of pathological events in the oral environment, resulting in inflammatory changes in gingival tissues, periodontal attachment losses, and the destruction of alveolar bone. The relationship between female sex hormones (estrogens) and periodontal disease in women during puberty, pregnancy, and menopause, as well as when taking oral contraceptives, has been widely reported (1,2).

Pregnancy is associated with remarkable changes in both endocrine function and immune reaction (3). Sheehan syndrome (SS), also known as postpartum pituitary necrosis, is a rare condition; it was first reported in 1937 by Sheehan et al. (4), who described a case of postpartum pituitary necrosis and insufficiency. The incidence of SS is 0.2 to 2.8 cases per 100,000 women in developed countries (5). This syndrome, which is caused by postpartum necrosis of the anterior lobe of the pituitary gland, can manifest clinically and pathologically in various ways, including agalactorrhea, amenorrhea, hypothyroidism, weakness, fatigue, dizziness, libido and weight loss, dry skin, pallor, hypoglycemia, adrenal crisis, and coma. The most common finding in SS is lactation that is difficult or fails to occur because of prolactin (PRL) deficiency. Furthermore, in many cases with SS, there is a sequential loss of somatotropic, gonadotropic, adrenocorticotrophic, and thyrotrophic functions (6). In this case report, we evaluate the periodontal status of a patient with SS who applied to our clinic because of periodontal problems.

## Case Report

The patient was a 63-year-old woman of 158 cm in height and 60 kg in weight; she had been diagnosed with SS in the endocrine department and referred to the periodontology clinic of Gülhane Faculty of Dentistry with complaints of gingival recessions, tooth loss, and gingival bleeding. In the intraoral examination of the patient, mobility was observed in some anterior teeth, with dense dental plaque accumulation and gums that were hyperemic and edematous. The patient declared that she was nonsmoking and that she brushed her teeth once a day but did not use an interface brush. First, medical and dental histories were taken. It was learned from the patient's medical history that she had had four children, and she was 18 when she initially gave birth; this baby had low birth weight. After the first birth, the patient had three more children who also had low birth weight. The patient told us that after her last birth, she was hospitalized for a while because of heavy bleeding. Following this heavy bleeding, the patient entered menopause at the age of 32. The patient lived for years without being diagnosed with SS,

but in 2018, she had to receive inpatient treatment at the Hacettepe University internal medicine clinic because of excessive fatigue, anorexia, and psychological depression. She was diagnosed with SS based on laboratory findings, pituitary magnetic resonance (MR) inspection, and her medical history.

After medical and dental histories were taken, extraoral and intraoral examinations were performed. In the patient's intraoral examination, it was observed that five teeth had been extracted, and there was migration and attachment loss in her teeth because of previous periodontal disease. Periodontal indexes, including the Plaque Index (PI), Gingival Index (GI), Bleeding on Probing (BOP), Periodontal Pocket Depth (PPD), Clinical Attachment Loss (CAL), and Recession Degree (RD), were recorded. The data containing all the patient's periodontal parameter values are given in Table 1. On panoramic radiography, it was revealed that there was a tooth root embedded in the bone in the posterior region of the left mandibula. It was observed in the radiographic examination of the patient that horizontal bone resorption occurred on the teeth, especially in the premolar region. From the clinical and radiological examination of the periodontal tissues, the patient was diagnosed with Stage IV, Grade C periodontitis (Figs. 1 and 2).




Figure 1. Patient's intraoral photographs.



Figure 2. Patient's panoramic radiography.

To control the patient's systemic condition, she was sent to the Endocrinology Clinic of Gülhane Medical Faculty, and a consultation was requested. As shown in the patient's laboratory results in Fig. 3, basal plasma levels of free T3 and T4, growth hormone (GH), PRL,

follicle stimulating hormone (FSH), luteinizing hormone (LH), progesterone, and total testosterone were significantly lower. In the pituitary MR examination, the pituitary gland was in the normal location, and the pituitary anterior gland height was found to be 1.1 mm, representing a significant decrease (Table 2).



**Figure 3.** MR image of the patient's pituitary gland.

Oral hygiene training was given to the patient after periodontal measurements. Scaling and root planning procedures were performed. The patient was called for control after 1 month and attended regular checkups at 3 and 6 months.

**Table 1.** Patient's clinical periodontal parameter value.

Periodontal Parameter Values		
Full mouth	PI	2.06
	GI	1.83
	BOP (%)	65.53
	PPD	2.31
	CAL	2.90
	RD	0.59

PI: Plaque Index, GI: Gingival Index, BOP: Bleeding on Probing, PPD: Periodontal Pocket Depth, CAL: Clinic Attachment Loss, RD: Recession Degree.

**Table 2.** Patient's laboratory results.

Laboratory Results		
Test	Results	Normal range
*Free T3	1.43	2.3-4.2
*Free T4	<0.40	0.78-1.48
Thyroid-Stimulating Hormone (TSH)	1.263	0.55-4.78
Estradiol	<10	<32.2(menopause)
*Follicle-Stimulating Hormone (FSH)	0.43	9.7-111 (menopause)
*Luteinising Hormone (LH)	0.38	15.9-54 (postmenopause)
*Prolactin	1.51	2.8-29.2
*Progesterone	<0.1	<0.73(postmenopause)
Parathormone (PTH)	36.4	15-65
*Growth Hormone	<0.05	<8
Adrenocorticotrophic hormone (ACTH)	12.53	0-46
25 Hydroxy Vitamin D	61.89	25-80
*Somatomedin-C	<15	63.4-223
*Total Testosterone	<0.13	0.14-0.76

## Discussion

The relationship between female sex hormones and periodontal disease in women during puberty, pregnancy, and menopause, as well as those taking oral contraceptives, has been widely reported (1,2). The presence of multiple hormonal deficiencies leads to the impairment of bone microarchitecture, which can cause osteopenia and even osteoporosis (7). Osteoporosis and periodontitis are both chronic diseases characterized by bone loss. Potential associations are of great clinical importance because of

the multifactorial etiology and common risk factors of these diseases. Indeed, recent studies have shown that there is a relationship between menopause, osteoporosis, alveolar bone resorption, and tooth loss (8,9). The progression of periodontal inflammation is regulated by systemic and oral environmental factors, including changes in physiological hormone levels during post-menopause (10).

In the past, postmenopausal osteoporosis was attributed solely to declining estrogen levels. In addition, estrogen replacement therapy was found to be effective in protecting against systemic bone loss and decreased alveolar bone loss (11). Reduction of

estrogen levels may affect the oral cavity epithelium, salivary gland function, and propensity for inflammatory processes related to periodontal tissues to emerge (12). Many researchers have demonstrated increased alveolar bone resorption involving estrogen deficiency (13,14).

PRL has been reported to be an important calcitropic hormone responsible for the provision of additional calcium during reproductive periods (15). One study demonstrated that at a lower PRL concentration of 10 ng/mL, PRL appeared to directly regulate human periodontal ligament fibroblast differentiation, as indicated by an upregulated mRNA expression of periost, the specific periodontal ligament differentiating marker involved in remodeling (16).

T3 and T4 are hormones secreted by the thyroid gland, and they have been shown to be fundamental to normal bone turnover (17). Thyroid dysfunctions are common health problems in the population and determining the influence of thyroid hormone imbalance in periodontitis may be important for the prevention of morbidity related to this condition when the association is present. The decrease in the levels of these hormones may promote a less competent immunogenic response to the infection induced by experimental periodontitis (18).

GH acts directly on the immune system, such as by priming monocytes for enhanced production of hydrogen peroxide and stimulating neutrophils to secrete superoxide anions associated with enhanced phagocytic activity (19). Therefore, deficiency in GH secretion by the pituitary gland or impaired function of its receptor could facilitate bacterial infections and cause tissue degradation, as occurs in periodontitis.

## Conclusions

Systemic diseases that affect hormonal conditions, such as SS, must be considered in the evaluation of oral health. Periodontitis that is exacerbated by or associated with systemic diseases is an important topic that must be considered and discussed in depth. Medical physicians should refer their patients to the dentist for consultation when evaluating patients' hormonal status and planning their treatment.

**Acknowledgments:** This study was presented as a full-text oral presentation at the 1<sup>st</sup> International Dental Research and Health Sciences Congress held between 20-22 May 2021.

**Patient Consent for Publication:** Written informed consent was obtained from the patient.

**Peer-review:** Externally peer-reviewed.

**Author Contributions:** Conception - S.C.Ş.; Design -N.I.S.; Supervision - A.O.K.; Materials - E.Ö.; Data Collection and/or Processing - S.C.Ş.; Analysis and/or Interpretation - N.I.S.; Literature Review - A.O.K.; Writer - E.Ö.; Critical Review - N.I.S.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study has received no financial support.

## References

1. Jafri Z, Bhardwaj A, Sawai M, Sultan N. Influence of female sex hormones on periodontium: A case series. *J Nat Sci Biol Med.* 2015; 6(Suppl 1):S146-9. ([Crossref](#))
2. Shapiro LF, Freeman K. The relationship between estrogen, estrogen receptors and periodontal disease in adult women: A review of the literature. *New York State Dental Journal*, 2014; 80(3), 30-34.
3. Amino N. et al. *Endocrine Journal* 2000; 47 (6), 645-655. ([Crossref](#))
4. Sheehan HL. Postpartum necrosis of the anterior pituitary. *J Pathol Bacteriol* 1937; 45:189-214. ([Crossref](#))
5. Kovacs K. Sheehan syndrome. *Lancet.* 2003; 361:520-2. ([Crossref](#))
6. Ozbey N, Inanc S, Aral F, Azezli A, Orhan Y, Sencer E, Molvalilar S. Clinical and laboratory evaluation of 40 patients with Sheehan's syndrome. *Isr J Med Sci* 1994; 30:826-829.
7. Gokalp D, Tuzcu A, Bahceci M, Arikan S, Ozmen CA, Cil T. Sheehan's syndrome and its impact on bone mineral density. *Gynecol Endocrinol.* 2009; 25:344-9. ([Crossref](#))
8. Oh TJ, Bashutski J, Giannobile WV. The interrelationship between osteoporosis and oral bone loss. *Grand Rounds Oral Syst Med* 2007; 2.
9. Guiglia R, Di Fede O, Lo Russo L, Sprini D, Rini GB, Campisi G. Osteoporosis, jawbones and periodontal disease. *Med Oral Patol Oral Cir Bucal.* 2013; 18(1):e93-9. ([Crossref](#))
10. Brennan RM, Genco RJ, Hovey KM, Trevisan M and Wactawski-Wende J. Clinical attachment loss, systemic bone density, and subgingival calculus in postmenopausal women. *J Periodontol* 2007; 78: 2104-2111. ([Crossref](#))
11. Duarte PM, Goncalves P, Casati MZ, de Toledo S, Sallum EA and Nociti FH Jr. Estrogen and alendronate therapies may prevent the influence of estrogen deficiency on the tooth-supporting alveolar bone: a histometric study in rats. *J Periodontol Res* 2006; 41: 541-546. ([Crossref](#))
12. Gupta R, Goel K, Solanki J. Gupta S. Oral manifestations of hypothyroidism: a case report. *J Clin Diagn Res.* 2014; 8:ZD20-22.
13. Kawamoto S, Nagaoka E. The effect of oestrogen deficiency on the alveolar bone resorption caused by traumatic occlusion. *J Oral Rehabil* 2000; 27:587-594. ([Crossref](#))
14. Kawamoto S, Ejiri S, Nagaoka E, Ozawa H. Effects of oestrogen deficiency on osteoclastogenesis in the rat periodontium. *Arch Oral Biol* 2002; 47:67-73. ([Crossref](#))
15. Charoenphandhu N, Wongdee K, Krishnamra N. Is prolactin the cardinal calcitropic maternal hormone? *Trends Endocrinol Metab.* 2010; 21:395-401. ([Crossref](#))
16. Surarit R, Krishnamra N, Seriwatanachai D. Prolactin receptor and osteogenic induction of prolactin in human periodontal ligament fibroblasts. *Cell Biol Int.* 2016; Apr;40(4):419-27. ([Crossref](#))
17. Bland R. Steroid hormone receptor expression and action in bone. *Clin Sci* 2000; 98:217-240. ([Crossref](#))
18. Bucay AH. Clinical hypothesis: application of AIDS vaccines together with thyroid hormones to increase their immunogenic effect. *Vaccine* 2007; 25: 6292-6293. ([Crossref](#))
19. Hattori N. Expression, regulation and biological actions of growth hormone (GH) and ghrelin in the immune system. *Growth Horm IGF Res* 2009. 19: 187-197. ([Crossref](#))